

Bactine Max Liquid Bandage
With Lidocaine

Drug Facts	
Active ingredients (w/w)	Purpose
Lidocaine 4%	External Analgesic
Uses ■ For temporary relief of pain or discomfort in minor cuts, scrapes and burns	
Warnings For external use only.	
Do not use ■ If you are allergic to any of the ingredients ■ in or near the eyes ■ over large areas of the body ■ longer than 1 week unless directed by a doctor	
Ask a doctor before use if you have ■ deep or puncture wounds ■ deep cuts ■ animal bites ■ serious burns	
When using this product ■ do not apply over raw surfaces or blistered areas	
Stop use and ask a doctor if ■ condition worsens ■ symptoms persist for more than 7 days, or clear up and occur again within a few days	
Keep out of reach of children. If swallowed, call a poison control center or get medical help right away.	
Directions ■ adults and children 2 years of age and older: ■ shake well ■ clean the affected area and apply a small amount up to 3 times daily and let dry ■ a second coating may be applied for extra protection ■ children under 2 years of age: ask doctor prior to use	
Other information Store at room temperature	
Inactive ingredients acrylates copolymer, alcohol denat, ammonium acrylates copolymer, benzalkonium chloride, disodium deceth-6 sulfosuccinate, disodium EDTA, laureth-30, propanediol, sodium dehydroacetate, water	
Questions? 1-844-241-5454 or Bactine.com	